

More on Meta-Stable Brane Configuration

Changhyun Ahn

Department of Physics, Kyungpook National University, Taegu 702-701, Korea

ahn@knu.ac.kr

Abstract

We describe the intersecting brane configuration of type IIA string theory corresponding to the meta-stable non-supersymmetric vacua in four dimensional $\mathcal{N} = 1$ supersymmetric $SU(N_c)$ gauge theory with an antisymmetric flavor, a conjugate symmetric flavor, eight fundamental flavors, m_f fundamental flavors and m_f antifundamental flavors. This is done by analyzing the $\mathcal{N} = 1$ supersymmetric $SU(2m_f - N_c + 4)$ magnetic gauge theory with dual matters and the corresponding dual superpotential.

1 Introduction

In the standard type IIA supersymmetric brane configuration of electric theory, the classical $\mathcal{N} = 1$ SQCD with gauge group $SU(N_c)$, N_f flavors of chiral superfields in the fundamental and antifundamental representations and vanishing superpotential can be described by N_c D4-branes suspended between NS5-brane and NS5'-brane along x^6 direction together with N_f D6-branes to the left of the NS5-brane, each of which is connected to the NS5-brane by a D4-brane. Similarly, in the standard type IIA supersymmetric brane configuration of magnetic theory, the classical $\mathcal{N} = 1$ SQCD with gauge group $SU(N_f - N_c)$, N_f flavors of chiral superfields in the fundamental and antifundamental representations and nonvanishing superpotential can be described by $(N_f - N_c)$ color D4-branes suspended between NS5'-brane and NS5-brane along x^6 direction together with N_f D6-branes to the left of the NS5'-brane, each of which is connected to the NS5'-brane by a flavor D4-brane [1, 2].

The dynamical supersymmetry breaking [3] can occur in $\mathcal{N} = 1$ SQCD with massive flavors where the supersymmetry is broken by rank condition through F-term equations. The geometric misalignment [4, 5, 6] of N_f flavor D4-branes connecting between NS5'-brane and D6-branes in the magnetic brane configuration can be interpreted as nontrivial F-term equations in the corresponding gauge theory. It turns out the minimal energy supersymmetry breaking IIA brane configuration can be described by coincident $(N_f - N_c)$ color D4-branes suspended between coincident $(N_f - N_c)$ D6-branes and NS5-brane along x^6 direction together with the remaining coincident N_c D6-branes to the left of the NS5'-brane, each of which is connected to the NS5'-brane by a flavor D4-brane [4, 5, 6]. For symplectic or orthogonal gauge groups, similar analysis can be done by adding orientifold 4-plane [5, 7].

The classical $\mathcal{N} = 1$ SQCD with gauge group $SU(N_c)$, N_f flavors of chiral superfields in the fundamental and antifundamental representations, a single adjoint field and nonvanishing superpotential can be described by N_c color D4-branes suspended between two coincident NS5-branes and NS5'-brane along x^6 direction together with N_f D6-branes at values of x^6 that are between those corresponding to the positions of the NS5-branes and NS5'-brane [1, 2]. The dynamical supersymmetry breaking [8, 9] can occur in $\mathcal{N} = 1$ SQCD with massive flavors plus a single adjoint by adding some mesonic deformation terms. It turns out the minimal energy supersymmetry breaking IIA brane configuration can be described by coincident $(2N_f - N_c)$ color D4-branes suspended between coincident $(2N_f - N_c)$ D6-branes and rotated two NS5-branes along x^6 direction, the coincident $(N_c - N_f)$ D6-branes to the left of the NS5'-brane, each of which is connected to the NS5'-brane by a flavor D4-brane, together with the remaining coincident $(N_c - N_f)$ D6-branes to the left of the NS5'-brane, each of which is connected to

the NS5'-brane by a tilted flavor D4-brane [10].

The classical $\mathcal{N} = 1$ SQCD with gauge group $SU(N_c)$, N_f flavors of chiral superfields in the fundamental and antifundamental representations, a symmetric flavor, a conjugate symmetric flavor and nonvanishing superpotential can be described by N_c color D4-branes suspended between rotated NS5-brane and rotated NS5'-brane along x^6 direction, an O6-plane, together with N_f D6-branes at values of x^6 that are between those corresponding to the positions of the rotated NS5-brane and NS5-brane (and their mirrors) [11, 2]. It turns out the minimal energy supersymmetry breaking IIA brane configuration can be described by coincident $(2N_f - N_c)$ color D4-branes suspended between coincident $(2N_f - N_c)$ D6-branes and NS5-brane along x^6 direction together with the remaining coincident $(N_c - N_f)$ D6-branes to the left of the NS5'-brane, each of which is connected to the NS5'-brane by a flavor D4-brane (and their mirrors) plus O6-plane [12].

In this paper, we study $\mathcal{N} = 1$ $SU(N_c)$ gauge theory with an antisymmetric flavor A , a conjugate symmetric flavor $\tilde{S}$, eight fundamental flavors $\hat{Q}$, m_f fundamental flavors Q and m_f antifundamental flavors $\tilde{Q}$ ¹ in the context of dynamical supersymmetric breaking vacua. The corresponding supersymmetric brane configuration in type IIA string theory was given by various groups in [13, 14, 15] sometime ago. See also [16]. In the related gauge theory analysis [17] alone, the turning on the superpotential $W = (A\tilde{S})^2$ was necessary to truncate the chiral ring and the dual description was given. From the brane analysis where, in general, two other quartic terms in the superpotential are present, this quartic superpotential can be obtained from the corresponding $\mathcal{N} = 2$ theory by integrating the massive adjoint field out.

For the limit of infinite mass for the adjoint field, the above quartic superpotential terms vanish because the coefficient function appears as an inverse of mass. Now we can add the mass term for quarks in the fundamental representation of the gauge group while keeping the infinite mass limit. Then we turn to the dual magnetic gauge theories by standard brane motions with appropriate linking number counting. It turns out that the dual magnetic theory giving rise to the meta-stable vacua is described by $\mathcal{N} = 1$ $SU(2m_f - N_c + 4)$ gauge theory with dual matter contents and the superpotential. Some of the F-term conditions cannot be satisfied by rank condition. Then the final nonsupersymmetric minimal energy brane configuration for this theory can be summarized in Figure 3².

In section 2, we review the type IIA brane configuration corresponding to the electric theory based on the $\mathcal{N} = 1$ $SU(N_c)$ gauge theory with above various matter contents. In

¹The proposal for the type IIA brane configuration describing the nonsupersymmetric meta-stable minimum in this theory is given by [5] already.

²There exists also other type of type IIA brane configuration [18] we do not discuss, describing the meta-stable vacuum in the $\mathbf{Z}_2$ orbifold of the conifold. See also other brane configuration in [19].

section 3, we construct the Seiberg dual magnetic theory which is $\mathcal{N} = 1$ $SU(3m_f - N_c + 4)$ gauge theory with corresponding dual matters as well as various gauge singlets, by brane motion and linking number counting. In section 4, we consider the infinite mass limit for the adjoint field of $\mathcal{N} = 2$ theory in order to obtain nonsupersymmetric meta-stable minimum where the gauge group is given by $SU(2m_f - N_c + 4)$ with matter contents and present the corresponding intersecting brane configuration of type IIA string theory. In section 5, we describe M-theory lift of the supersymmetry breaking type IIA brane configuration we have considered in section 4. Finally, in section 6, we give the summary of type IIA brane configuration corresponding to meta-stable vacua of various gauge theories.

2 The $\mathcal{N} = 1$ supersymmetric electric brane configuration

Let us describe $\mathcal{N} = 1$ $SU(N_c)$ gauge theory with an antisymmetric flavor A , a conjugate symmetric flavor $\tilde{S}$, eight fundamental flavors $\hat{Q}$, m_f fundamental flavors Q and m_f antifundamental flavors $\tilde{Q}$ [13, 14, 15]. The anomaly-free global symmetry group determined by the gauge symmetry on the D6-branes is given by $SU(m_f)_L \times SO(8)_L \times SU(m_f)_R$ with three $U(1)$ 's³. The type IIA brane configuration consists of three NS5-branes having different x^6 values, N_c D4-branes suspended between them, $2m_f$ D6-branes, eight “half” D6-branes extending all the way from $x^7 = 0$ to $x^7 = \infty$, an orientifold 6 plane($O6^+$ -plane) located at negative x^7 carrying +4 RR charge and an orientifold 6 plane($O6^-$ -plane) located at positive x^7 carrying -4 RR charge⁴.

Let us introduce two complex coordinates, as usual,

$$v \equiv x^4 + ix^5, \quad w \equiv x^8 + ix^9.$$

According to $\mathbf{Z}_2$ symmetry of $O6$ -plane (0123789) sitting at $v = 0$ and $x^6 = 0$, the coordinates (x^4, x^5, x^6) transform as $-(x^4, x^5, x^6)$. Among three NS5-branes, the middle NS5'-brane (012389) which is located at $x^7 = 0$ is stuck on an $O6$ -plane and the two outer NS5-branes are

³Note that this gauge theory looks similar to [17] in the sense that the matter contents are the same but the global symmetries and the superpotential are different. Over there, the global symmetry is $SU(m_f + 8)_L \times SU(m_f)_R$ with three $U(1)$'s and the superpotential has more simple form.

⁴The orientifold action identifies the two factors of product gauge group $SU(N_c) \times SU(N_c)$ and projects the bifundamental representation onto one antisymmetric tensor and one conjugate symmetric tensor since there exists different projection acting on the different sides of the D4-branes [14]. Moreover, the matter coming from the half D6-branes exists. The brane configuration of $\mathcal{N} = 1$ supersymmetric gauge theory with $SU(N_L) \times SU(N_R)$ and matter in the bifundamental and fundamental representations was found in [20, 21, 22].

placed in a $\mathbf{Z}_2$ symmetric manner from $\mathcal{N} = 2$ brane configuration as in Figure 1. The left NS5-brane is rotated by an angle θ in (v, w) plane (denoted by $NS5_\theta$ -brane) while the right NS5-brane is rotated by an angle $-\theta$ in the same plane (denoted by $NS5_{-\theta}$ -brane). Our convention for this angle is different from the one used in [13]. Similarly, the left m_f D6-branes are rotated by an angle ω in (v, w) plane independently and denote them as $D6_\omega$ -branes and the right m_f D6-branes appear as $D6_{-\omega}$ -branes in a $\mathbf{Z}_2$ symmetric manner⁵. The middle NS5'-brane divides the O6-plane into two disconnected regions corresponding to the positive and negative x^7 , denoted by $O6^-$ -plane and $O6^+$ -plane respectively. The RR charge of the O6-plane jumps from $+4$ (in negative region of x^7) to -4 (in positive region of x^7) as we cross the NS5'-brane [23]. The part of $O6^-$ -plane has more eight semi-infinite (or half) D6-branes (0123789) embedded in it. That is, four half D6-branes (and their mirrors) are present. This is necessary to preserve the charge conservation, or to cancel the discontinuity of the RR charge [24, 13, 14, 15].

For this brane setup, the classical superpotential in electric theory is given by the quartic terms, flavor symmetry breaking term and a mass term for quarks as follows⁶

$$W = -\frac{1}{2\mu} \left[(A\tilde{S})^2 + Q\tilde{S}A\tilde{Q} + (Q\tilde{Q})^2 \right] + \hat{Q}\tilde{S}\hat{Q} + mQ\tilde{Q}, \quad \mu \equiv \tan\left(\frac{\pi}{2} - \theta\right) \quad (2.1)$$

where A is an antisymmetric field, $\tilde{S}$ is a conjugate symmetric field and $\hat{Q}$ is the 8 fundamental fields as well as m_f fundamental fields Q and m_f antifundamental fields $\tilde{Q}$ and the adjoint mass μ is related to the rotation angle θ as above [25, 2].

Recall that starting with the brane configuration with $\mathcal{N} = 2$ supersymmetry, one can rotate one or both of the NS5-branes or one or more of the D6-branes such that the $\mathcal{N} = 1$ supersymmetry is preserved. The mass of the adjoint chiral superfield breaks the $\mathcal{N} = 2$ supersymmetry to $\mathcal{N} = 1$ supersymmetry. Keeping one of the NS5-branes and all the D6-branes fixed and rotating the other NS5-brane corresponds to changing the mass of the adjoint chiral superfield. On the other hand, rotating both NS5-branes relative to the D6-branes keeping the two NS5-branes and all the D6-branes parallel corresponds to changing the coupling between the adjoint chiral superfield and quarks [2].

When $NS5_\theta$ -brane is parallel to $D6_\omega$ -branes or $\theta = \omega$ (and their mirrors also), then the last two terms in the quartic terms above (2.1) are not present because the coefficient function

⁵There exists the figure showing the various angles in the product gauge group. One can make the following identifications explicitly : for the left branes $\theta = \frac{\pi}{2} - \theta_1$ and $\omega = \frac{\pi}{2} - \theta_1 - \omega_1$ where θ_1 and ω_1 are defined in the figure 5 of [14].

⁶If there exist k coincident $NS5_\theta$ -branes (and their mirrors also), then the superpotential takes the form as $W = (\mu\Phi^{k+1} + A\Phi\tilde{S} + Q\Phi\tilde{Q}) + \hat{Q}\tilde{S}\hat{Q} + mQ\tilde{Q}$, in general. For the particular case where $NS5_\theta$ -brane is parallel to the $D6_\omega$ -branes (and their mirror also) indicating that there is no $Q\Phi\tilde{Q}$ term, one obtains $W = \frac{1}{2\mu}(A\tilde{S})^{k+1} + \hat{Q}\tilde{S}\hat{Q} + mQ\tilde{Q}$ [14].

$\sin(\theta - \omega)$ appears in front of the $Q\Phi\tilde{Q}$ before integrating out the adjoint field Φ ⁷. We did not insert the dependence on this coefficient function into (2.1) explicitly in order to make it simple and actually in the limit of $\mu \rightarrow \infty$ (or $\theta \rightarrow 0$), all the quartic terms vanish. The field theory analysis [17]⁸ corresponding to the superpotential $(A\tilde{S})^2$ which was necessary to truncate the chiral ring provides the magnetic dual description. For the infinite limit of μ , it is evident that the superpotential becomes $\hat{Q}\tilde{S}\hat{Q} + mQ\tilde{Q}$. Since the eight half D6-branes are sitting on the top of O6-plane, the flavor symmetry breaking term $\hat{Q}\tilde{S}\hat{Q}$ in the superpotential indicates that the flavor symmetry $SU(m_f + 8)_L$ is broken to $SU(m_f)_L \times SO(8)_L$.

Then, the $\mathcal{N} = 1$ supersymmetric electric brane configuration in type IIA string theory realized in the gauge theory we study can be summarized as follows:

- Middle NS5'-brane with worldvolume (012389) with $v = 0 = x^6 = x^7$ ⁹.
- Left $NS5_\theta$ -brane with worldvolume by both (0123) and two spatial dimensions in (v, w) plane and with negative x^6 .
- Right $NS5_{-\theta}$ -brane with worldvolume by both (0123) and two spatial dimensions in (v, w) plane and with positive x^6 .
- Left m_f D6 $_\omega$ -branes with worldvolume by both (01237) and two spatial dimensions in (v, w) plane and with negative x^6 . Before the rotation, the distance from D4-branes in the v direction is nonzero.
- Right m_f D6 $_{-\omega}$ -branes with worldvolume by both (01237) and two spatial dimensions in (v, w) plane and with positive x^6 .
- Eight semi-infinite D6-branes with worldvolume (0123789) with $x^7 \geq 0$ and $x^6 = 0$. The

⁷When the $2m_f$ D6-branes approach and are placed on top of the O6-plane, the tree level superpotential is given by $(\mu\Phi^2 + A\Phi\tilde{S} + Q\Phi\tilde{Q}) + (Q\tilde{S}Q + \tilde{Q}A\tilde{Q}) + \hat{Q}\tilde{S}\hat{Q} + mQ\tilde{Q}$ where the global symmetry is enhanced to $SO(2m_f + 8)_L \times Sp(m_f)_R$. Note that the matter contents in Q and $\tilde{Q}$ are doubled [13, 14, 15]. That is, an antisymmetric flavor A , a conjugate symmetric flavor $\tilde{S}$, eight fundamental flavors $\hat{Q}$, $2m_f$ fundamental flavors Q and $2m_f$ antifundamental flavors $\tilde{Q}$. Instead, if we put $2m_f$ D6'-branes (0123457) approaching to the O6-plane, then the global symmetry becomes $SO(2m_f)_L \times SO(8)_L \times Sp(m_f)_R$. According to the result of [15, 11], the dual magnetic gauge group from brane configuration is given by $SU(2m_f - N_c + 4)$ and these extra two Φ -independent terms in the superpotential, $Q\tilde{S}Q + \tilde{Q}A\tilde{Q}$, can be interpreted as mesonic perturbation in the magnetic theory.

⁸When the baryonic operator $\tilde{B}_n = \tilde{S}^n \tilde{Q}^{N_c - n} \tilde{Q}^{N_c - n}$ gets an expectation value, then the initial gauge group $SU(N_c)$ along this flat direction is broken to $SO(n)$ with antisymmetric tensor $\hat{A}$ and $2m_f$ vectors as well as the singlets and the superpotential will be $\hat{A}^2$ [13]. Along this flat direction, the original theory flows to other theory in the IR [26]. Similarly, when the baryonic operator $B_n = A^n Q^{N_c - 2n}$ gets an expectation value, then the initial gauge group $SU(N_c)$ is broken to $Sp(n)$ with a conjugate symmetric tensor $\hat{\tilde{S}}$ and $2m_f$ flavors and the superpotential will be $\hat{\tilde{S}}^2$.

⁹According to $\mathbf{Z}_2$ symmetry of O6-plane, there exist two possible orientations of a middle NS5-brane. It should be its own mirror under this action. It can extend in the v direction given in [12] or it can extend in w direction. The second case provides the chiral theory [13].

four half D6-branes are mirror to the remaining four half D6-branes with respect to O6-plane¹⁰.

- Semi-infinite $O6^+$ -plane with worldvolume (0123789) with $v = 0 = x^6$ and $x^7 \leq 0$.
- Semi-infinite $O6^-$ -plane with worldvolume (0123789) with $v = 0 = x^6$ and $x^7 \geq 0$.
- N_c D4-branes with worldvolume (01236) with $v = 0 = w$.

We draw the type IIA electric brane configuration in Figure 1 which was basically given in [13, 14, 2] already but the only difference is to put $D6_{\pm\omega}$ -branes in the nonzero v direction in order to obtain nonzero masses for the quarks which are necessary to obtain the meta-stable vacua. Also one can imagine the brane configuration for $\theta = 0$ (in which the two outer $NS5_{\pm\theta}$ become two NS5-branes) from Figure 1.




Figure 1: The $\mathcal{N} = 1$ supersymmetric electric brane configuration for the $SU(N_c)$ gauge theory with $A, \tilde{S}$ (that are strings stretching between D4-branes located at the left hand side of O6-plane and those at the right hand side of O6-plane), $\hat{Q}$ (that are strings stretching between D6-branes which are on top of $O6^-$ -plane and D4-branes) and $Q, \tilde{Q}$ (that are strings stretching between D6-branes and D4-branes). The O6-plane is divided into half by the middle NS5'-brane in the x^7 -direction. The origin of the coordinates (x^6, v, w) is located at the intersection of NS5'/O6/D6-branes (a combination of a middle NS5'-brane, $O6^+$ -plane, $O6^-$ -plane and eight half D6-branes above) and D4-branes.

¹⁰Our convention for the location of these eight fundamentals are different from the one in [2] because the axis of x^7 is reversed.

3 The $\mathcal{N} = 1$ supersymmetric magnetic brane configuration

In order to obtain the magnetic dual theory in the context of brane configuration, we move the D6-branes and NS5-branes through each other [2] and use the linking numbers [27] for the computation of D4-branes which are created during this process. In other words, the magnetic theory in the brane picture is obtained by interchanging D6-branes and NS5-branes with their mirror images while preserving the linking number.

The linking number [27] for the D6-branes is given by $L_6 = \frac{1}{2}(n_{5L} - n_{5R}) + n_{4R} - n_{4L}$ where $n_{5L,R}$ are the NS5-branes to the left or right of the D6-branes and $n_{4L,R}$ are the D4-branes to the left or right of the D6-branes. After we move the left $D6_\omega$ -branes to the right all the way (and their mirrors, right $D6_\omega$ -branes to the left) past the NS5'-brane and $NS5_\theta$ -brane, the linking number L_6 of a $D6_\omega$ -brane becomes $L_6 = 1 - n_{4L}$, which should be equal to the original $L_6 = -1$ before the brane motion, according to the conservation of linking number. Then the n_{4L} becomes 2 and we must add $2m_f$ D4-branes to the left side of all m_f $D6_\omega$ -branes (and their mirrors). See Figure 2 for the creation of these D4-branes.

Next, we move the left $NS5_\theta$ -brane to the right all the way past O6-plane (and its mirror, right $NS5_\theta$ -brane to the left), the linking number of $NS5_\theta$ -brane, $L_5 = \frac{m_f}{2} + \frac{1}{2}(4) - N + 2m_f$ using the formula $L_5 = \frac{1}{2}(n_{6L} - n_{6R}) + n_{4R} - n_{4L}$ where $n_{6L,R}$ are the D6-branes to the left or right of the NS5-branes, $n_{4L,R}$ are the D4-branes to the left or right of the NS5-branes. Originally, it was $L_5 = -\frac{m_f}{2} - \frac{1}{2}(4) + N_c$ from Figure 1 before the brane motion. Therefore, we are left with the number of colors [13] in the magnetic theory

$$N = 3m_f - N_c + 4.$$

The magnetic dual theory ¹¹ has gauge group $SU(N)$ where $N = 3m_f - N_c + 4$ ¹² and a conjugate symmetric $\tilde{s}$ and an antisymmetric a chiral multiplet together with m_f fundamental q , 8 fundamental $\hat{q}$ and m_f antifundamental $\tilde{q}$ chiral multiplets and moreover there exist gauge singlet fields

$$M_0 \equiv Q\tilde{Q}, \quad M_1 \equiv Q\tilde{S}A\tilde{Q}, \quad \tilde{M} \equiv \hat{Q}\tilde{Q} \quad (3.1)$$

¹¹In [17], the number of colors is equal to $N = 3m_f - 12 - N_c$ for m_f fundamentals, $(m_f - 8)$ antifundamentals, an antisymmetric tensor and a conjugate symmetric tensor. By shifting as $m_f \rightarrow m_f + 8$ in order to compare with the gauge theory we consider and match with the matter contents, this becomes $N = 3m_f + 12 - N_c$. The two theories between the one in [13, 14] and the one in [17] are connected to each other. By giving the masses to all eight $\hat{Q}$'s in the electric theory, the former theory will reduce to the latter theory.

¹²When k $NS5_\theta$ -branes (and its mirrors) are present, as in the notation of footnote 6, then by doing successive Seiberg dual, the magnetic dual gauge group becomes $SU((2k+1)m_f - N_c + 4(2k-1))$ where $2km_f$ term comes from k coincident $NS5_{\pm\theta}$ -branes, m_f term comes from a middle NS5'-brane and $4(2k-1)$ constant term is due to the O6-plane.

as well as other type of gauge singlets $P \equiv Q\tilde{S}Q$ and $\tilde{P} \equiv \tilde{Q}A\tilde{Q}$ that are symmetric and antisymmetric in their flavor indices respectively ¹³. Among newly created $2m_f$ D4-branes during the process from the electric theory to magnetic theory, the m_f D4-branes (when m_f $D6_\omega$ -branes cross a middle NS5'-brane) in the magnetic theory can split on the $D6_{-\omega}$ -branes in the most general way and a total of m_f^2 massless modes correspond to the m_f^2 components of the magnetic meson M_1 . On the other hand, the remaining m_f D4-branes (when m_f $D6_\omega$ -branes cross the right NS5 $_{-\theta}$ -brane) in the magnetic theory can split similarly and m_f^2 massless modes correspond to the same components of the magnetic meson M_0 . In the electric theory, if we move the $D6_\omega$ -branes on top of O6-plane, as in the footnote 7, we have seen the terms like as $P \equiv Q\tilde{S}Q$ and $\tilde{P} \equiv \tilde{Q}A\tilde{Q}$. Here the fundamental chiral multiplets Q of the gauge group originate from 4-6 strings connecting D4-branes to $D6_{\pm\omega}$ -branes ending a middle NS5'-brane from positive x^7 while the antifundamental chiral multiplets $\tilde{Q}$ of the gauge group come from 4-6 strings connecting D4-branes to $D6_{\pm\omega}$ -branes ending a middle NS5'-brane from negative x^7 [15, 13, 14, 2]. A meson $\tilde{M}$ corresponds to 4-4 strings stretching between the right D4-branes and the left D4-branes.

The magnetic superpotential for arbitrary θ and ω is given by

$$W_{dual} = (a\tilde{s})^2 + M_1 q\tilde{q} + M_0 q\tilde{s}a\tilde{q} + Pq\tilde{s}q + \tilde{P}\tilde{q}a\tilde{q} + (\hat{q}\tilde{s}\hat{q} + \tilde{M}\hat{q}\tilde{q}) + mM_0$$

where the first five terms are obtained from the field theory analysis [17], the next two terms are due to the presence of flavor symmetry breaking (2.1) in the electric theory and the last term comes from the mass of quarks in the electric theory ¹⁴. In particular, P and $\tilde{P}$ -terms arise when the m_f $D6_\omega$ -branes intersect with its mirror m_f $D6_{-\omega}$ -branes around the location of a middle NS5'-brane. In next section, by considering the subset of this process from the electric theory to magnetic theory, we'll show how the brane configuration for the meta-stable supersymmetry breaking states arises.

We draw the type IIA magnetic brane configuration in Figure 2 which was given in [13, 14, 2] already but the only difference is to put $D6_{\pm\omega}$ -branes in the nonzero v direction in order to obtain nonzero masses for the quarks which are necessary to obtain the meta-stable vacua. Compared with the previous Figure 1, this Figure 2 looks similar to Figure 1 but there exist

¹³ For the brane picture in the footnote 7, after we move the left NS5 $_\theta$ -brane to the right all the way past O6-plane (and its mirror, right NS5 $_{-\theta}$ -brane to the left), the linking number of NS5 $_\theta$ -brane is given by $L_5 = \frac{(2m_f)}{2} + \frac{(4)}{2} - N$. Note that due to the doubling of the flavors of Q and $\tilde{Q}$, the number of D6-branes are $2m_f$ on top of O6-plane. Originally, it was $L_5 = -\frac{(2m_f)}{2} - \frac{(4)}{2} + N_c$ before the brane motion. Then the magnetic gauge group one finds is $SU(2m_f - N_c + 4)$ as in footnote 7.

¹⁴ As for the brane configuration in footnote 7, by adding the composite operators $Q\tilde{S}Q$ and $\tilde{Q}A\tilde{Q}$ in electric theory, one gets the generalized magnetic superpotential by adding P - and $\tilde{P}$ -terms into W_{dual} , observed in [15].

$2m_f$ D4-branes(and its mirrors) which will be reduced to m_f D4-branes(and its mirrors) for infinite adjoint mass limit in next section. Also one can imagine the brane configuration for $\theta = 0$ (in which the two outer $NS5_{\pm\theta}$ become two NS5-branes) from Figure 2.




Figure 2: The $\mathcal{N} = 1$ supersymmetric magnetic brane configuration for the $SU(N = 3m_f - N_c + 4)$ gauge theory with $a, \tilde{s}, \hat{q}, q$ and $\tilde{q}$. The $2m_f$ D4-branes connecting between $NS5_{-\theta}$ -brane and $D6_{-\omega}$ -branes are the dual gauge singlets corresponding to the mesons M_0 and M_1 (3.1). As will be explained in section 4, in the $\mu \rightarrow \infty$ limit (or $\theta \rightarrow 0$ limit), only m_f D4-branes connecting NS5-brane and D6-branes appear(and their mirrors). In the superpotential, it contains either M_0 or M_1 -dependent term depending on which way the brane motion occurs.

4 Nonsupersymmetric meta-stable brane configuration

When the D6-branes are parallel to a middle NS5'-brane or perpendicular to NS5-branes in the electric theory(in other words, the angles are given by $\omega = \frac{\pi}{2}$ and $\theta = 0$) in the $\mu \rightarrow \infty$ limit¹⁵, let us describe the nonsupersymmetric meta-stable brane configuration¹⁶. First rotate $D6_{\omega}$ -branes to the v direction and make them to be parallel to a middle NS5'-brane(and its mirrors also). After we move the left $D6_{\omega=\frac{\pi}{2}}$ -branes to the right all the way(and their mirrors, right $D6_{-\omega=-\frac{\pi}{2}}$ -branes to the left) past the middle NS5'-brane and the right NS5-brane(= $NS5_{-\theta=0}$ -brane), the linking number L_6 of a single D6-brane becomes $L_6 = \frac{1}{2} - n_{4L}$,

¹⁵In this case, the D4-branes can move freely in v direction along the two outer NS5-branes and A and $\tilde{S}$ are free to get vacuum expectation values [17, 14].

¹⁶For the case of $\theta = 0 = \omega$ (when the D6-branes are perpendicular to a middle NS5'-brane), the superpotential contains M_1 -dependent term because the m_f D4-branes created during the Seiberg dual process correspond to the singlet M_1 , as in [12].

which should be equal to the original linking number $L_6 = -\frac{1}{2}$. Let us denote $D6_{\omega=\pm\frac{\pi}{2}}$ -branes by unrotated D6-branes (0123789). Then we should add m_f D4-branes, corresponding to the meson M_0 , to the left side of all the right m_f D6-branes (and their mirrors) in Figure 2. Note that the difference, compared with previous case where $0 < \theta \leq \frac{\pi}{2}$, appears here when a D6-brane crosses the middle NS5'-brane. Due to the parallelness of these (D6-branes and NS5'-brane), there is no creation of D4-branes. This fact will affect the number of color D4-branes below.

Next, we move the left NS5-brane(=NS5 $_{\theta=0}$ -brane) to the right all the way past O6-plane (and its mirror, right NS5-brane to the left), and then the linking number of NS5-brane becomes $L_5 = \frac{m_f}{2} + \frac{1}{2}(4) - N + m_f$. Note that since the NS5-brane is nonparallel to the O6-plane (actually they are perpendicular to each other), one can see the contribution from the O6-plane which plays the role of four D6-branes, contrary to the result of [12] in which the outer NS5'-branes were parallel to an O6-plane. Originally, the linking number was $L_5 = -\frac{m_f}{2} - \frac{1}{2}(4) + N_c$. This implies that the number of D4-branes in magnetic theory, N , becomes

$$N = 2m_f - N_c + 4$$

which is the same as the case for the theory of the footnote 7.

The magnetic superpotential corresponding to the electric superpotential $W = \hat{Q}\tilde{S}\hat{Q} + mQ\tilde{Q}$ with $\mu \rightarrow \infty$ is given by

$$W_{dual} = M_0 q \tilde{s} a \tilde{q} + m M_0 + (\tilde{q} \tilde{s} \hat{q} + \tilde{M} \hat{q} \tilde{q}). \quad (4.1)$$

When we take $\mu \rightarrow \infty$, the quartic terms from (2.1) vanish and this leads to the fact that there is no such term like $(a\tilde{s})^2$ in the magnetic superpotential we are considering. Our particular route for brane motion, as we observed before, does not allow for us to have M_1 term which also prohibits the presence of $M_1 q \tilde{q}$. Moreover since the D6-branes are parallel to the middle NS5'-brane ($\omega = \frac{\pi}{2}$), there are no P or $\tilde{P}$ dependent terms. Eventually, we are left with the result of (4.1).

Here q and $\tilde{q}$ are fundamental and antifundamental for the gauge group indices and anti-fundamentals for the flavor indices. The fields a and $\tilde{s}$ are antisymmetric tensor and conjugate symmetric tensor for the gauge group indices respectively with no flavor indices. Then, $q\tilde{s}a\tilde{q}$ has rank N while m has a rank m_f . Therefore, the F-term condition, the derivative the superpotential W_{dual} with respect to M_0 , cannot be satisfied if the rank m_f exceeds N . This is so-called rank condition. Other F-term equations are satisfied by taking the vacuum expectation values of $\tilde{M}$ and $\hat{q}$ to vanish.

The classical moduli space of vacua can be obtained from F-term equations. From the F-terms F_q and $F_{\tilde{s}}$, one gets $\tilde{s}a\tilde{q}M_0 = 0 = a\tilde{q}M_0q + \hat{q}\hat{q}$. Similarly, one obtains $\tilde{q}M_0q\tilde{s} = 0 = M_0q\tilde{s}a + \tilde{M}\hat{q}$ from the F-terms F_a and $F_{\tilde{q}}$. Moreover, there is a relation $q\tilde{s}a\tilde{q} + m = 0$ from the F-term F_{M_0} . From the conditions $a\tilde{q}M_0 = 0 = M_0q\tilde{s}$ and $\tilde{M} = 0 = \hat{q}$, which satisfy the above four equations for nonzero vacuum expectation values $q, \tilde{q}, a$ and $\tilde{s}$, one can fix the form of M_0 and part of $q\tilde{s}$ and $a\tilde{q}$, which can be fixed by using $F_{M_0} = 0$ further, and one obtains the following solutions

$$\langle q\tilde{s} \rangle = \begin{pmatrix} \sqrt{m}e^\phi \mathbf{1}_N \\ 0 \end{pmatrix}, \langle a\tilde{q} \rangle = \begin{pmatrix} \sqrt{m}e^{-\phi} \mathbf{1}_N & 0 \end{pmatrix}, \langle M_0 \rangle = \begin{pmatrix} 0 & 0 \\ 0 & \Phi_0 \mathbf{1}_{m_f-N} \end{pmatrix} \quad (4.2)$$

as well as $\langle \tilde{M} \rangle = 0 = \langle \hat{q} \rangle$ where $\Phi_0 \mathbf{1}_{m_f-N}$ is an arbitrary $(m_f - N) \times (m_f - N)$ matrix and the zeros of $\langle q\tilde{s} \rangle$ and $\langle a\tilde{q} \rangle$ are $(m_f - N) \times N$ and $N \times (m_f - N)$ zero matrices respectively. Similarly, the zeros of $m_f \times m_f$ matrix M_0 are assumed also. Then Φ_0 and $(\sqrt{m}e^\phi, \sqrt{m}e^{-\phi})$ parametrize a pseudo-moduli space. Let us expand around on a point on (4.2), as done in [3]. That is,

$$q = \begin{pmatrix} q_0 \mathbf{1}_N + \frac{1}{\sqrt{2}}(\delta\chi_+ + \delta\chi_-) \mathbf{1}_N \\ \delta\varphi \end{pmatrix}, \tilde{s} = (\tilde{s}_0 + \delta\tilde{X}) \mathbf{1}_N, a = (a_0 + \delta X) \sigma_2 \otimes \mathbf{1}_{\frac{N}{2}},$$

$$\tilde{q} = \begin{pmatrix} \tilde{q}_0 \sigma_2 \otimes \mathbf{1}_{\frac{N}{2}} + \frac{1}{\sqrt{2}}(\delta\chi_+ - \delta\chi_-) \sigma_2 \otimes \mathbf{1}_{\frac{N}{2}} & \delta\tilde{\varphi} \sigma_2 \otimes \mathbf{1}_{\frac{m_f-N}{2}} \end{pmatrix}, M_0 = \begin{pmatrix} \delta Y & \delta Z \\ \delta\tilde{Z} & \Phi_0 \mathbf{1}_{m_f-N} \end{pmatrix}$$

together with the variations of $\tilde{M}$ and $\hat{q}$ that are $m_f \times 8$ and $8 \times N$ matrices respectively. Then the superpotential (4.1) becomes

$$W_{dual}^{fluct} = \Phi_0 (\delta\varphi \tilde{s}_0 a_0 \delta\tilde{\varphi} + m) + \delta Z \delta\varphi \tilde{s}_0 a_0 \tilde{q}_0 + \delta\tilde{Z} q_0 \tilde{s}_0 a_0 \delta\tilde{\varphi} \\ + \left(\frac{1}{\sqrt{2}} \delta Y \delta\chi_+ \tilde{s}_0 a_0 \tilde{q}_0 + \dots \right) + (\text{cubic}) \quad (4.3)$$

where (cubic) stands for the terms that are cubic or higher in the fluctuations and $\dots$ contains some parts from the last two terms in (4.1) that are not relevant. Then to quadratic order, the model splits into two sectors where the first piece given by the first line of (4.3) is an O'Raifeartaigh type model and the second piece given by the second line of (4.3) is supersymmetric and will not contribute to the supertrace. The fields $\delta\chi_\pm, \delta Y, \delta X$ and $\delta\tilde{X}$ couple to the supersymmetry breaking fields $\delta\varphi$ and $\delta\tilde{\varphi}$ via terms of cubic and higher order in the fluctuations. Then, the remaining relevant terms of superpotential are given by

$$W_{dual}^{rel} = \Phi_0 (\delta\hat{\varphi} \delta\hat{\tilde{\varphi}} + m) + \delta Z \delta\hat{\varphi} a_0 \tilde{q}_0 + \delta\tilde{Z} q_0 \tilde{s}_0 \delta\hat{\tilde{\varphi}}$$

where $\delta\hat{\varphi} \equiv \delta\varphi \tilde{s}_0$ and $\delta\hat{\tilde{\varphi}} \equiv a_0 \delta\tilde{\varphi}$. At one loop, the effective potential $V_{eff}^{(1)}$ for Φ_0 can be obtained from this relevant part of superpotential which consists of the matrix M and N of

[28] where the defining function $\mathcal{F}(v^2)$ can be computed. Using the equation (2.14) of [28] of $m_{\Phi_0}^2$ and $\mathcal{F}(v^2)$, one gets that $m_{\Phi_0}^2$ will contain $(\log 4 - 1) > 0$, by taking the limit where $q_0 \tilde{s}_0 \rightarrow \sqrt{m} e^\phi$ and $a_0 \tilde{q}_0 \rightarrow \sqrt{m} e^{-\phi}$, as in (4.2). This implies that these vacua are stable. In the brane configuration from Figure 2, since the N D4-branes can slide along the two NS5-branes when $\theta = 0$, the vacuum expectation values of a and $\tilde{s}$ can be 2×2 block-diagonalized.

By recombination of N of D4-branes connecting D6-branes and NS5-brane with those connecting NS5'-brane and NS5-brane and moving them in v direction (and their mirrors) from Figure 2, the minimal energy supersymmetry breaking brane configuration is shown in Figure 3 that was observed also in [5]. This phenomenon occurs all the examples found in [4, 5, 6, 10, 7, 12]. Compared with the brane configuration of [12], the locations of NS5-brane and NS5'-brane are interchanged each other. Roughly speaking, when the NS5-brane moves along the x^6 direction and two NS5'-branes move to the origin from the Figure 3 of [12], then one obtains the Figure 3 of this paper. Undoing the Seiberg dual in the context of [5]. Some entries in the dual quarks $q, \tilde{q}$ and dual tensors $a, \tilde{s}$ acquire the nonzero expectation values in terms of the eigenvalues of mass matrix m to minimize the F-term F_{M_0} in this dual gauge theory. This corresponds exactly to the geometric recombination and movement along the v direction above. When one gets the nonzero vacuum expectation values for $\langle M_0 \rangle, \langle q \rangle$, and $\langle \tilde{q} \rangle$, then the separation of the D4-branes along the middle NS5'-brane corresponds to the mass of two index tensor $(a\tilde{s})$.

5 M-theory description of nonsupersymmetric metastable brane configuration

The M5-brane lives in (0123) directions and wraps on a Riemann surface inside (4568910) directions. The Taub-NUT space (45610) is parametrized by two complex variables (v, y) while flat two-dimensions (89) are by a complex variable w . One of the complex structures of Taub-NUT space (45610) can be described by embedding it in a three complex dimensional space with coordinates (x, t, v) . The M5-brane curve corresponding to the type IIA brane configuration shown in Figure 1, in a background space [29] of $xt = (-1)^{m_f} v^4 \prod_{k=1}^{m_f} (v^2 - e_k^2)$ where e_k is the position of the D6-branes in the v direction, can be characterized by a cubic equation in t [11, 12]. For fixed x , the coordinate t corresponds to y and for fixed y , the coordinate x corresponds to $1/t$.

At nonzero string coupling $g_s \neq 0$, the two kinds of NS5-branes can bend due to their interactions with the D4-branes and D6-branes. Let us consider the asymptotic behavior such that the rotated curve should have at $w \rightarrow \infty$ and $w = 0$. One can read off the




Figure 3: The nonsupersymmetric minimal energy brane configuration for the $SU(N = 2m_f - N_c + 4)$ gauge theory with $a, \tilde{s}, q, \tilde{q}$ as well as $\hat{q}$. This can be obtained from the magnetic description of Figure 2 by putting $\theta = 0$ and $\omega = \frac{\pi}{2}$ and with the condition that the number of D4-branes connecting between D6-branes and NS5'-brane are reduced to a half. The $(m_f - N)$ D4-branes connecting between NS5'-brane and D6-branes can move freely along w direction and this leads to nonzero vacuum expectation values of the meson M_0 (3.1). In section 5, we move each half of the middle NS5'-brane to the $\pm v$ direction holding everything else fixed, instead of moving D6-branes.

behaviors of the left, middle, right NS5-branes respectively as in [12]. Then the behavior of the supersymmetric M5-brane curves can be written as follows by realizing that the role of NS5-brane and NS5'-brane (the behaviors of the coordinates v and w) are exchanged into each other:

1. $w \rightarrow \infty$ limit implies

$$v \rightarrow \pm m, \quad y \sim w^{m_f+2} + \dots \quad \text{NS' asymptotic region.}$$

2. $v \rightarrow \infty$ limit implies

$$w \rightarrow 0, \quad y \sim v^{2m_f - N_c + 4} + \dots \quad \text{NS}_L \text{ asymptotic region,}$$

$$w \rightarrow 0, \quad y \sim v^{N_c} + \dots \quad \text{NS}_R \text{ asymptotic region.}$$

Along the line of [6, 7, 12], each half of the central NS5'-brane is moved to the $\pm v$ direction holding everything else fixed, instead of moving D6-branes (and their mirrors). In this process, the corresponding mirrors are placed in appropriate way. Since the nonsupersymmetric brane configuration in Figure 3 implies that only the NS5'-brane is deformed by turning on the mass for the quarks, only NS5'-brane is nonholomorphic. The remaining NS5-brane and D6-branes remain unchanged. The ansatz for this nonholomorphic curve corresponding to the NS5'-brane with boundary condition above can be made similarly.

The harmonic function appearing in the Taub-NUT space, sourced by the the left coincident m_f D6-branes located at $x^6 = 0$, $O6^+$ -plane(which is equivalent to 4 D6-branes) located at $x^6 = l$, and the right coincident m_f D6-branes located at $x^6 = 2l$, can be written as explicitly, by putting the right charges and the locations of them,

$$V(s) = 1 + \frac{m_f R}{\sqrt{f(s)^2 + s^2}} + \frac{4R}{\sqrt{f(s)^2 + (s-l)^2}} + \frac{m_f R}{\sqrt{f(s)^2 + (s-2l)^2}}. \quad (5.1)$$

Note that when we compare with the usual $\mathcal{N} = 1$ SQCD with massive flavors developed in [6], the last two terms in (5.1) coming from the effect of O6-plane and mirrors of D6-branes in our gauge theory are present newly. When half of the middle NS5'-brane is located at $v = \frac{\Delta x}{\ell_s^2}$ from Figure 3 (and its mirror appears at opposite value of v) and $f(s) = \Delta x$ ¹⁷, then there exists the equation (A.3) of [6] which is valid for arbitrary form for the harmonic function. Then, this first order differential equation leads to the solution for $g(s)$,

$$g(s) \sim \left(s - l + \sqrt{(\Delta x)^2 + (s-l)^2} \right)^{\frac{2}{N_c}} \prod_{j=0}^1 \left[s - 2lj + \sqrt{(\Delta x)^2 + (s-2lj)^2} \right]^{\frac{m_f}{2N_c}}$$

where the s -independent integration constant is fixed by the boundary condition above classification 2. The function $g(s)$ does not vanish even if Δx vanishes. Then w that is proportional to $g(s)$ does not vanish and y is not equal to zero. This is not consistent with the statement that $y = 0$ only if $v = 0$. Therefore, the extra piece in the potential (5.1) doesn't remove the instability from a new M5-brane mode occurring at some point during the continuation to M-theory description of SQCD.

6 Summary and outlook

Let us make some comments on the type IIA brane configuration corresponding to the meta-stable vacua for different kinds of gauge theories.

- Orthogonal group with a symmetric tensor flavor and massive vectors

As observed in [12], it is hard to imagine how the meta-stable vacua supersymmetry breaking brane configuration arises because there is no extra NS5-brane we need to have. The type IIA supersymmetry breaking brane configuration of meta-stable vacua corresponding to $SU(2N_f - N_c)$ gauge theory with matters is shown in Figure 3 of [12]. If we go on baryonic branch(i.e., we move a middle NS5-brane to x^7 direction), the baryon gets an expectation value along this flat direction and the gauge group is broken to $SO(n)$ with a symmetric

¹⁷Remember that the ansatz for the nonholomorphic curve implies that $x^4 = f(s)$ and $x^5 = 0$ and for the mirror of NS5'-brane we should have $f(s) = -\Delta x$ due to the $\mathbf{Z}_2$ symmetry of O6-plane.

tensor as well as vectors. The superpotential has a quadratic expression on this symmetric tensor because the D4-branes will cross the $O6^+$ -plane in the region giving a projection to the orthogonal gauge group [14]. Therefore, the type IIA supersymmetry breaking brane configuration is the Figure 3 of [12] with a shift of NS5-brane to x^7 direction. The gauge theory analysis on this fact was given in [17] or the flow from mezzanine to orchestra in the IR of A_k model [26] corresponds to this example in the gauge theory side. In order to obtain higher order term for the superpotential in terms of the symmetric tensor, we need to insert a multiple of outer NS5'-branes.

- Symplectic group with an antisymmetric tensor flavor and massive fundamental flavors

From the Figure 3 of [12] where $O6^+$ -plane is replaced by $O6^-$ -plane, if we move a middle NS5-brane to x^7 direction, the baryon gets an expectation value along this flat direction. Then the gauge group is broken to $Sp(n)$ with an antisymmetric tensor flavor as well as fundamental flavors and the superpotential has a quadratic expression on this antisymmetric tensor because the D4-branes will cross the $O6^-$ -plane in the region which gives a projection to the symplectic gauge group. Similarly, the type IIA supersymmetry breaking brane configuration is the Figure 3 of [12] where $O6^+$ -plane is replaced by $O6^-$ -plane with a shift of NS5-brane to x^7 direction.

- Orthogonal group with an adjoint(antisymmetric tensor) and massive vectors

From Figure 3 above in this paper, we can move NS5'-brane to $+x^7$ direction. The expectation value on this operator in magnetic dual breaks the dual gauge group $SU(N)$ to $SO(n)$ with an adjoint as well as vectors and a quadratic superpotential for this adjoint because the D4-branes will cross the $O6^+$ -plane in the region which gives a projection to the orthogonal gauge group [13, 14]. Therefore, the type IIA supersymmetry breaking brane configuration is the Figure 3 in this paper with a shift of NS5-brane to $+x^7$ direction. The gauge theory analysis on this fact was given in [17] or the flow from balcony to mezzanine in the IR of A_k model [26] corresponds to this example, in the gauge theory side.

On the other hand, according to [30], this theory can be realized in terms of $O4^+$ -plane, NS5-brane, NS5'-brane, D4-branes and D6-branes. In principle, one can follow the method of [8, 9] by adding the corresponding mesonic deformations and one expects that one can obtain the meta-stable vacua to this theory in the gauge theory side.

- Symplectic group with an adjoint(symmetric tensor) and massive fundamental flavors

From Figure 3, we can move NS5'-brane to $-x^7$ direction. The expectation value on this operator in magnetic dual breaks the dual gauge group $SU(N)$ to $Sp(n)$ with an adjoint and a quadratic superpotential for this adjoint because the D4-branes will cross the $O6^-$ -plane in the region which gives a projection to the symplectic gauge group [13, 14]. Therefore, the

type IIA supersymmetry breaking brane configuration is the Figure 3 in this paper with a shift of NS5-brane to $-x^7$ direction.

Similarly, one realizes that this gauge theory corresponds to the brane configuration which consists of $O4^-$ -plane, NS5-brane, NS5'-brane, D4-branes and D6-branes [30]. It is evident that one also obtains the corresponding brane configuration from Figure 4 of [10] by inserting all the mirrors with respect to $O4^-$ -plane appropriately.

Then it is natural to ask what happens for the product gauge groups. For example, in the gauge theory side [17], the theory of $SU(N_1) \times SO(N_2)$ with X , which transform as fundamental in $SU(N_1)$ and vector in $SO(N_2)$, a conjugate symmetric tensor $\tilde{S}$ in $SU(N_1)$, as well as fundamentals for $SU(N_1)$ and vectors for $SO(N_2)$ can confine either $SU(N_1)$ factor or $SO(N_2)$ factor. For the latter, the IR duality is given by an unitary gauge theory with a symmetric tensor flavor and fundamental flavors. See also relevant paper [31] where the matter contents are different.

According to the result of [32], by rotating $\mathcal{N} = 2$ brane configuration, a single smooth supersymmetric M5-brane description gives rise to the monopole and meson expectation values which are matching with the field theory computation through confining phase superpotential [33]. See also the relevant discussions for symplectic and orthogonal gauge groups in [34, 35]. The $\mathcal{N} = 1$ supersymmetric magnetic configuration for this case where k' NS5'-branes are connected by $(2N_f - N_c)$ D4-branes to a single NS5-brane on their right and by a single D4-brane to each of N_f D6-branes on their left was given in [30]. For $k' = 2$ case, the gauge theory analysis was given in [36] where the magnetic dual superpotential corresponding to the electric superpotential $W = X^3$ with an adjoint X is given by $W_{dual} = Y^3 + M_1 \tilde{q} Y q + M_2 \tilde{q} q$ with a dual adjoint Y , dual quarks $q, \tilde{q}$ and gauge singlets $M_1 \equiv Q \tilde{Q}$ and $M_2 \equiv Q X \tilde{Q}$ for electric quarks $Q, \tilde{Q}$ and an adjoint X . It would be interesting to study whether there exist nonsupersymmetric meta-stable vacua.

Acknowledgments

I would like to thank A. Hanany and K. Landsteiner for discussions. This work was supported by grant No. R01-2006-000-10965-0 from the Basic Research Program of the Korea Science & Engineering Foundation. I thank KIAS(Korea Institute for Advanced Study) for hospitality where this work was undertaken.

References

- [1] S. Elitzur, A. Giveon and D. Kutasov, Phys. Lett. B **400**, 269 (1997)

- [2] A. Giveon and D. Kutasov, *Rev. Mod. Phys.* **71**, 983 (1999)
- [3] K. Intriligator, N. Seiberg and D. Shih, *JHEP* **0604**, 021 (2006)
- [4] H. Ooguri and Y. Ookouchi, *Phys. Lett. B* **641**, 323 (2006)
- [5] S. Franco, I. Garcia-Etxebarria and A. M. Uranga, *JHEP* **0701**, 085 (2007)
- [6] I. Bena, E. Gorbatov, S. Hellerman, N. Seiberg and D. Shih, *JHEP* **0611**, 088 (2006)
- [7] C. Ahn, *Phys. Lett. B* **647**, 493 (2007)
- [8] A. Amariti, L. Girardello and A. Mariotti, *JHEP* **0612**, 058 (2006)
- [9] A. Amariti, L. Girardello and A. Mariotti, [arXiv:hep-th/0701121].
- [10] C. Ahn, *Class. Quant. Grav.* **24**, 1359 (2007)
- [11] K. Landsteiner, E. Lopez and D. A. Lowe, *JHEP* **9807**, 011 (1998)
- [12] C. Ahn, [arXiv:hep-th/0701145].
- [13] K. Landsteiner, E. Lopez and D. A. Lowe, *JHEP* **9802**, 007 (1998)
- [14] I. Brunner, A. Hanany, A. Karch and D. Lust, *Nucl. Phys. B* **528**, 197 (1998)
- [15] S. Elitzur, A. Giveon, D. Kutasov and D. Tsabar, *Nucl. Phys. B* **524**, 251 (1998)
- [16] K. Landsteiner and E. Lopez, *Nucl. Phys. B* **516**, 273 (1998)
- [17] K. A. Intriligator, R. G. Leigh and M. J. Strassler, *Nucl. Phys. B* **456**, 567 (1995)
- [18] R. Argurio, M. Bertolini, S. Franco and S. Kachru, *JHEP* **0701**, 083 (2007)
- [19] R. Kitano, H. Ooguri and Y. Ookouchi, *Phys. Rev. D* **75**, 045022 (2007)
- [20] J. H. Brodie and A. Hanany, *Nucl. Phys. B* **506**, 157 (1997)
- [21] A. Giveon and O. Pelc, *Nucl. Phys. B* **512**, 103 (1998)
- [22] J. Erlich, A. Naqvi and L. Randall, *Phys. Rev. D* **58**, 046002 (1998)
- [23] N. J. Evans, C. V. Johnson and A. D. Shapere, *Nucl. Phys. B* **505**, 251 (1997)
- [24] A. Hanany and A. Zaffaroni, *Nucl. Phys. B* **509**, 145 (1998)

- [25] J. L. F. Barbon, Phys. Lett. B **402**, 59 (1997)
- [26] J. H. Brodie and M. J. Strassler, Nucl. Phys. B **524**, 224 (1998)
- [27] A. Hanany and E. Witten, Nucl. Phys. B **492**, 152 (1997)
- [28] D. Shih, [arXiv:hep-th/0703196].
- [29] E. Witten, Nucl. Phys. B **500**, 3 (1997)
- [30] S. Elitzur, A. Giveon, D. Kutasov, E. Rabinovici and A. Schwimmer, Nucl. Phys. B **505**, 202 (1997)
- [31] E. Lopez and B. Ormsby, JHEP **9811**, 020 (1998)
- [32] J. de Boer and Y. Oz, Nucl. Phys. B **511**, 155 (1998)
- [33] S. Elitzur, A. Forge, A. Giveon, K. A. Intriligator and E. Rabinovici, Phys. Lett. B **379**, 121 (1996)
- [34] C. Ahn, Phys. Lett. B **426**, 306 (1998)
- [35] C. Ahn, K. Oh and R. Tatar, J. Geom. Phys. **28**, 163 (1998)
- [36] O. Aharony, J. Sonnenschein and S. Yankielowicz, Nucl. Phys. B **449**, 509 (1995)